

Geometric phase as a determinant of a qubit– environment coupling

**Quantum Information
Processing**

ISSN 1570-0755

Volume 10


Number 1

Quantum Inf Process (2010)

10:85-96

DOI 10.1007/s11128-010-0178-

x



Geometric phase as a determinant of a qubit– environment coupling

J. Dajka · J. Łuczka · P. Hänggi

Received: 20 January 2010 / Accepted: 20 April 2010 / Published online: 5 May 2010
© Springer Science+Business Media, LLC 2010

Abstract We investigate the qubit geometric phase and its properties in dependence on the mechanism for decoherence of a qubit weakly coupled to its environment. We consider two sources of decoherence: dephasing coupling (without exchange of energy with environment) and dissipative coupling (with exchange of energy). Reduced dynamics of the qubit is studied in terms of the rigorous Davies Markovian quantum master equation, both at zero and non–zero temperature. For pure dephasing coupling, the geometric phase varies monotonically with respect to the polar angle (in the Bloch sphere representation) parameterizing an initial state of the qubit. Moreover, it is antisymmetric about some points on the geometric phase–polar angle plane. This is in distinct contrast to the case of dissipative coupling for which the variation of the geometric phase with respect to the polar angle typically is non-monotonic, displaying local extrema and is not antisymmetric. Sensitivity of the geometric phase to details of the decoherence source can make it a tool for testing the nature of the qubit–environment interaction.

Keywords Geometric phase · Dephasing · Dissipation · Open system · Davies theory

1 Introduction

One of the key obstructions of an effective implementation of quantum algorithms is related to the ubiquitous problem of decoherence in real quantum objects [1]. Quantum

J. Dajka (✉) · J. Łuczka
Institute of Physics, University of Silesia, 40-007 Katowice, Poland
e-mail: dajka@phys.us.edu.pl

P. Hänggi
Institute of Physics, University of Augsburg, 86135 Augsburg, Germany

decoherence is generic as it results from the imperfect isolation of the quantum system from its environment. Decoherence can be diminished under very special conditions such as e.g. the presence of the decoherence free subspaces [2,3] or via the application of tailored, external control schemes [4]. A promising novel direction in quantum information relates to so called holonomic or topological quantum computations [5,6] allowing for a substantial reduction of decoherence [7,8]. The essence of this method consists in encoding the information in the holonomy related to the geometric phase of the quantum evolution [9,10]. The geometric phase can be expressed as a path integral and via the Stokes theorem, can be converted into a surface integral. Therefore, it behaves like a geometric area. A quantity like an area is less dependent on the details of time evolution and therefore is less affected by changes of environmental conditions or an imperfect control, and hence, is typically more robust. This is the key attribute that makes geometric phases attractive for the implementations of fault-tolerant quantum computation. Some suggestions have been presented to realize this objective, e.g. in NMR experiments [7], ion traps [11], neutral atoms in cavity QED [12], quantum dots [13] or Josephson junction devices [14,15]. The performance of holonomic quantum gates under various conditions has been studied recently [16,17].

The quantum evolution in the presence of decoherence is generically non-unitary. Therefore, the notion of geometric phase needs to be extended. There are several extensions of the geometric phase concept for systems which are either in a mixed state or/and undergo a non-unitary evolution. The first attempt towards this goal is given in Uhlmann [18], being rather of mathematical character. The other are based on quantum trajectories [19,20], quantum interferometry [21–25] and the state purification (kinematic approach) [26,27]. For non-unitary quantum evolution there is no commonly accepted scheme of defining the geometric phase in open quantum systems [28–32]. Here we use the approach based on state purification as proposed in Ref. [27]. This so defined geometric phase has been extensively studied in various contexts [33–37]. One of the appealing ‘advantages’ of studying the phase defined in Tong et al. [27] is that it can be measured with a carefully prepared interferometric experiment [21–25,27]. Our reasoning is thus guided by its potential for experimental implementation.

There is no unique method of describing the time evolution of open quantum systems and there are several schemes to treat such systems which however typically give rise to non-equivalent dynamics [38,39]. One scheme consists in the derivation of a reduced system dynamics, via tracing over the degrees of freedom of the environment. Except some few exactly solvable models [38,40,41] it is not clear how to relate the reduced dynamics to the microscopically first principles dynamics based upon the Hamiltonian structure of quantum dynamics [38,42]. The exactly solvable model of pure dephasing has been applied for studying quantum channels [43] or in exploring the dynamics of quantum entanglement [44–48]. Despite its simplicity the highly non-trivial properties of geometric phase of qubits has also been discussed [49–52]. One of the most successful examples of constructing reduced dynamics is the Davies approximation scheme [53]. Within this approach, starting with the general ‘system-bath-interaction’ Hamiltonian, one obtains, in a mathematically rigorous way, a Markovian master equation form of a quantum system weakly coupled to the environment which preserves positivity and yields the correct equilibrium Gibbs state

[54,55]. This approach has been applied to various problems of statistical physics, quantum optics, solid state physics and quantum information, e.g. for studying entanglement dynamics in bipartite systems [56].

In this paper we apply the Davies master equation to study the geometric phase of the qubit coupled to the bosonic bath. Various families of a coupling and different coupling strength are shown to result in a qualitative and quantitative modification of the geometric phase. This behavior could suggest a method to resolve the nature of the qubit–bath coupling: In particular, the dephasing coupling presents not only a mere theoretical construction but can be realized in experiments within tailored regimes [57]. In order to keep this study self-contained, we briefly review the notion of the geometric phase for a non-unitarily evolving qubit and then present the qubit master equation derived from the Davies theory.

2 Geometric phase

Generally, the time evolution of the qubit reduced density matrix $\rho(t)$ is neither unitary nor Markovian [38,42]. It is constructed as the mapping

$$\rho(t) = \Lambda(t, t_0)\rho(t_0) \tag{1}$$

obeying some properties depending on the specific circumstances and approximations such as e.g. the celebrated complete positivity condition [58]. In order to exploit the approach to the geometric phase based on state purification [27] we have to present the reduced density matrix Eq.(1) in the spectral-decomposition form

$$\rho(t) = \sum_{i=1}^2 p_i(t)|w_i(t)\rangle\langle w_i(t)|, \tag{2}$$

where $p_i(t)$ and $|w_i(t)\rangle$ are the eigenvalues and the eigenvectors of the matrix $\rho(t)$, respectively. The geometric phase $\Phi(t)$ associated with such an evolution is defined as follows [27]:

$$\begin{aligned} \Phi(t) = \arg & \left[\sum_{i=1}^2 [p_i(0)p_i(t)]^{1/2} \langle w_i(0)|w_i(t) \rangle \right. \\ & \left. \times \exp \left(- \int_0^t \langle w_i(s)|\dot{w}_i(s) \rangle ds \right) \right], \end{aligned} \tag{3}$$

where \arg denotes the argument of the complex number, $\langle w_i|w_j \rangle$ is a scalar product and the dot indicates the derivative with respect to time s . For convenience, we assume the initial time being $t_0 = 0$. For the sake of completeness we sketch here, following Ref. [27], the derivation of Eq. (3). The mixed state defined by the density matrix (2) can be lifted to a pure state $|\Psi(t)\rangle$ in a larger Hilbert space, i.e.,

$$|\Psi(t)\rangle = \sum_{i=1}^2 \sqrt{p_i(t)} |w_i(t)\rangle \otimes |a_i\rangle, \tag{4}$$

where the vectors $|a_i\rangle$ span the Hilbert space of an arbitrary ancilla. This is known as a purification of the density matrix $\rho(t)$ in the sense that $\rho(t)$ is a partial trace of the density matrix $|\Psi(t)\rangle\langle\Psi(t)|$ over the ancilla Hilbert space. With the time evolution of the purified system one can associate the ‘Pancharatnam’ relative phase

$$\alpha(t) = \arg\langle\Psi(0)|\Psi(t)\rangle \tag{5}$$

which contains both the gauge-dependent part (a dynamical phase) and a gauge-independent part. The central result of Tong et al. [27] is to extract from Eq. (5), by a proper choice of the ‘parallel transport condition’, the purification-independent part which can be termed a geometric phase because it is gauge invariant and reduces to the known results in the limit of an unitary evolution [59,60]. The final result is then given by Eq. (3).

As mentioned in the Introduction, this phase—contrary to other attempts of extending the notion of geometric phase for a non-unitary evolving quantum system—has a direct physical meaning as it can be measured via interferometric experiments [27], i.e. one can construct the purification of the quantum system such that the relative phase Eq. (5) reduces to the geometric phase Eq. (3) after suitably defined ‘compensating unitary’ cutting of the dynamical part of the relative phase [27].

3 Weak coupling regime of qubit reduced dynamics

The evolution operator $\Lambda(t, t_0)$ defined by Eq.(1) or its infinitesimal generator \mathcal{L} defined by the equation

$$\frac{d}{dt}\rho(t) = \mathcal{L}\{\rho(t)\} \tag{6}$$

can be obtained in a few cases only; namely for stylized, exactly solvable models or in the limiting regimes such as the weak coupling limit or the singular coupling limit [58]. We consider a qubit coupled to a bosonic environment at temperature T . The Hamiltonian of such a system is chosen in the form [38]:

$$H = H_Q \otimes \mathcal{I} + \mathcal{I} \otimes H_B + H_I \otimes V_B, \tag{7}$$

$$H_B = \int_0^\infty E(k) a^\dagger(k) a(k) dk, \tag{8}$$

$$V_B = \int_0^\infty g(k) [a^\dagger(k) + a(k)] dk. \tag{9}$$

The operators $a^\dagger(k)$ and $a(k)$ denote the creation and annihilation boson operators, respectively. The qubit Hamiltonian and the interaction are assumed to take the form

$$H_Q = \frac{\varepsilon}{2}\sigma_z, \quad H_I = \hbar\mu_x\sigma_x + \hbar\mu_z\sigma_z, \tag{10}$$

where σ_i are the Pauli operators, ε is the qubit energy splitting and the dimensionless parameters μ_x and μ_z are coupling constants. Let us remark that if $\mu_x \equiv 0$ the qubit energy operator H_Q is an integral of motion, i.e. it commutes with the total Hamiltonian H leaving the expectation value of the corresponding energy observable unchanged. This situation defines the well known exactly solvable model of pure dephasing [40,41]. A non-vanishing μ_x is then characteristic for exchange of energy and related dissipation processes.

For an uncorrelated initial state $\rho(0) \otimes w(\beta)$ taken as a product of an arbitrary qubit density matrix $\rho(0)$ and the equilibrium Gibbs state of the environment $w(\beta) = \exp(-\beta H_B)/\text{Tr}[\exp(-\beta H_B)]$ with $\beta = 1/k_B T$ (k_B is the Boltzmann constant), the Davies approximation for the Markovian kernel yields the following Markovian master equation [53–55]

$$\frac{d}{dt}\rho(t) = \mathcal{L}_H\{\rho(t)\} + \mathcal{L}_R\{\rho(t)\}, \tag{11}$$

where the ‘conservative’ and ‘dissipative’ parts read as follows

$$\mathcal{L}_H\{\rho(t)\} = -\frac{i}{\hbar} \left[\left(H_Q + \sum_{k,l=-1}^1 \hbar s(\Omega_{kl}) A_{kl}^\dagger A_{kl} \right), \rho(t) \right], \tag{12}$$

$$\begin{aligned} \mathcal{L}_R\{\rho(t)\} = & \frac{1}{2} \sum_{k,l=-1}^1 c(\Omega_{kl}) \\ & \times \left([A_{kl}\rho(t), A_{kl}^\dagger] + [A_{kl}, \rho(t)A_{kl}^\dagger] \right), \end{aligned} \tag{13}$$

where, see in Refs. [56,58],

$$\begin{aligned} A_{kl} &= \mathcal{P}_k H_I \mathcal{P}_l, \quad \mathcal{P}_{\pm 1} = |\pm 1\rangle\langle\pm 1|, \\ \Omega_{kl} &= (\lambda_k - \lambda_l)/\hbar, \quad \lambda_{\pm 1} = \pm\varepsilon/2. \end{aligned} \tag{14}$$

The states $|1\rangle$ and $|-1\rangle$ denote the excited state and ground state of the qubit, respectively. The quantity $c(\omega)$ is the Fourier transform of the autocorrelation function of the bath operator V_B calculated in the Gibbs state $w(\beta)$ of the bath, namely,

$$c(\omega) = \int_{-\infty}^{\infty} e^{-i\omega t} \text{Tr} \left[w(\beta) V_B e^{itH_B/\hbar} V_B e^{-itH_B/\hbar} \right] dt \tag{15}$$

and its Hilbert transform defines the function $s(\omega)$ in the following way

$$s(\omega) = \frac{P}{2\pi} \int_{-\infty}^{\infty} \frac{c(x)}{x - \omega} dx, \tag{16}$$

where P indicates the Cauchy principal value of the integral.

In order to treat the complex qubit–environment interaction encoded in $g(k)$ in Eq. (9) it is convenient to introduce the spectral density

$$D(\omega) = \int dk |g(k)|^2 \delta(\omega(k) - \omega). \tag{17}$$

We further limit our consideration to the strictly Ohmic environment for which this spectral density is linear with respect to ω for small frequencies and exhibits an exponential cut-off frequency ω_c , thereby exhibiting no non-physical ultraviolet divergences. Explicitly, this spectral density reads

$$D(\omega) = \frac{\alpha}{2} \omega \exp(-\omega/\omega_c), \tag{18}$$

where the dimensionless parameter α characterizes the strength of the environmental influence on the qubit.

Within this choice [56,58]

$$c(\omega) = \frac{\pi\alpha}{2} \left(|\omega| \frac{\exp(\beta\hbar|\omega|) + 1}{\exp(\beta\hbar|\omega|) - 1} + \omega \right) \exp(-|\omega|/\omega_c) \tag{19}$$

and $s(\omega)$ is determined via the relation in Eq. (16).

In principle one can solve Eq. (11) using the Bloch vector formalism to obtain the coupled evolution equations for mean values $\langle \sigma_k(t) \rangle$, $k = x, y, z$ to obtain the reduced density matrix as $\rho(t) = (1/2)[1 + \langle \sigma_x(t) \rangle \sigma_x + \langle \sigma_y(t) \rangle \sigma_y + \langle \sigma_z(t) \rangle \sigma_z]$. This form allows to extract the spectral decomposition Eq. (2) and the phase $\Phi(t)$. Such an explicit form of the geometric phase result is, however, rather cumbersome without exhibiting much physical insight. We thus refrain from presenting such analytical details, but present here the full analysis of the geometric phase by numerical means.

4 Analysis of geometric phase

From Eqs. (1)–(3) it follows that in order to determine the geometric phase at arbitrary time $t > 0$, we must specify the initial state of the qubit. We consider the following class of initial states

$$|\theta\rangle = \cos(\theta/2)|1\rangle + \sin(\theta/2)|-1\rangle, \tag{20}$$

where θ is the polar angle in the Bloch sphere representation. The corresponding initial statistical operator $\rho(0)$ takes the form

$$\rho(0) = |\theta\rangle\langle\theta|. \tag{21}$$

One of the eigenvalues of this operator is zero, say $p_2(0) = 0$ in Eq. (3), and it does not contribute to the geometric phase. This simplifies Eq. (3) in that only one term of the sum survives. The evolution of the freely evolving qubit, with $\mu_x = \mu_z = 0$ in Eq. (10), is cyclic with the time-period $\mathcal{T} = 2\pi\hbar/\varepsilon$ and it acquires the geometric phase [60]

$$\Phi_0 = \pi[1 + \cos(\theta)], \quad \text{mod}(2\pi), \tag{22}$$

which can serve as a reference for studying the influence of the environment. In the case of a coupling to an environment, the evolution of the qubit is not cyclic any longer. However, below we consider the phase $\Phi = \Phi(\mathcal{T})$ after the time $\mathcal{T} = 2\pi\hbar/\varepsilon$ in order to study the role of coupling to the environment and for comparison with Eq. (22).

The simplest situation occurs for pure dephasing; i.e. when $\mu_x = 0$ so that the qubit energy does not change. The results presented in Fig. 1 show that the geometric phase plotted as a function of the initial state of the qubit (i.e. as a function of the parameter θ in Eq. (20)) approaches zero (modulo 2π) with increasing coupling strength μ_z . We observe that it varies drastically in the regime near $\theta = \pi/2$ and varies weakly outside this region. Moreover, the phase vanishes for $\theta \rightarrow 0$ (i.e. for the initially excited state $|1\rangle$) and $\theta \rightarrow \pi$ (i.e. for the ground state $|-1\rangle$). This finding corroborates the results for the phase in the exactly solvable model of pure dephasing with arbitrary (not only weak) coupling [49–52]. For the presentation as in Fig. 1, we note that the function $\Phi(\theta)$ is antisymmetric about the point $\{\theta, \Phi\} = \{\pi/2, \pi\}$, i.e. the relation

$$\Phi\left(\frac{\pi}{2} + \theta\right) = 2\pi - \Phi\left(\frac{\pi}{2} - \theta\right) \quad \text{for } \theta \in [0, \pi/2] \tag{23}$$

holds. It can be interpreted as a rotation symmetry around the point $\{\theta, \Phi\} = \{\pi/2, \pi\}$.

A most intriguing behavior on the role of the environment emerges when $\mu_x \neq 0$; i.e. when the qubit–environment interaction is allowed to exchange energy with the

Fig. 1 (Color online) Dependence of the geometric phase $\Phi = \Phi(\mathcal{T})$ on the initial state of the qubit which is parameterized by the angle θ on the Bloch sphere. The qubit is coupled to a purely dephasing Ohmic environment, i.e. $\mu_x = 0$ in Eq. (10). The remaining parameters are: $\alpha = 10^{-2}$, $\hbar\omega_c/\varepsilon = 10^2$ and $T = 0$





Fig. 2 (Color online) Geometric phase $\Phi = \Phi(T)$ versus initial polar angle θ for selected values of the dissipative qubit–Ohmic environment coupling strength μ_x . The dephasing coupling strength is set at $\mu_z = 0$ in Eq. (10), and the remaining parameters are the same as in Fig. 1



qubit system. The results depicted in Fig. 2 show the qualitative changes in the geometric phase properties for increasing dissipation coupling strength μ_x as a function of the polar angle θ . Note that the geometric phase Φ in Fig. 2 is plotted differently from Fig. 1 with Φ/π varying within the interval $(-1, 1)$. We have decided to make this change in order to avoid confusing jump-like behavior of $\Phi(\theta)$ in vicinity of the polar angle $\theta = \pi/2$. E.g. the curve corresponding to the case $\mu_x = 0.05$ in Fig. 2 is very similar to the curve corresponding to the case $\mu_z = 0.1$ in Fig. 1. However, presented in the $\Phi/\pi \in (-1, 1)$ interval it exhibits jump-like behavior which is an artefact of the way the plot is done. Let us recall again here that the phase Φ is defined modulo 2π .

For small values of μ_x , the geometric phase is close to that for the isolated qubit, cf. $\mu_x = 0.05$ in Fig. 2 when compared with Fig. 1 but with Φ varying there $\Phi/\pi \in (0, 2)$. When μ_x increases the function $\Phi(\theta)$ exhibits a local maximum and minimum, see the case $\mu_x = 0.3$ in Fig. 2. For larger value of the coupling strength μ_x (the case $\mu_x = 0.4$) the geometric phase is an increasing function of the polar angle θ till to the value $\theta = \pi/2$ reaching a local maximum. Next, it decreases as $\theta \rightarrow \pi$. In comparison to the dephasing coupling, in this case we can find at least three distinguishing features of the geometric phase. Firstly, we note breaking of antisymmetry of $\Phi(\theta)$, being in distinct contrast to the case of pure dephasing ($\mu_x = 0$), cf. Fig. 1. Secondly, the dependence of the phase on the initial state parameterized by θ is non-monotonic, exhibiting a local maximum and a minimum. Thirdly, the geometric phase Φ vanishes for $\theta \rightarrow \pi$ (i.e. for the ground state) but not necessary so for $\theta \rightarrow 0$ (i.e. for the excited state).

One can observe that for a fixed μ_x , the dephasing process controlled by μ_z does not change the qualitative properties of the geometric phase Φ , see Figs. 1 and 3. Pure dephasing affects only the off-diagonal elements of the density matrix, becoming closer to the maximally mixed qubit state. The geometric phase in a quantum evolution of such states vanishes. In a general energy-exchanging process the time dependence of the density matrix is more complex and the geometric phase Φ seemingly quantifies this fact. Moreover, the stability of geometric phase with respect to decoherence is crucial for effectiveness of holonomic quantum computation [5, 13]. It is evident that the stability of phase can be significantly improved via a proper choice of the initial state determined by θ in Eq. (20).





Fig. 3 (Color online) Role of a qubit–environment coupling on the geometric phase $\Phi = \Phi(T)$ versus initial state preparation θ for a qubit that is coupled to an Ohmic environment. The dissipative coupling strength is set at $\mu_x = 0.3$. The influence of dephasing is depicted for four different coupling strengths μ_z . The remaining parameters are the same as in Fig. 1

Fig. 4 (Color online) The influence of varying temperature T on the geometric phase $\Phi(T)$ versus initial state preparation θ is depicted for a dissipative qubit with $\mu_x = 0.3$ and zero dephasing, i.e. $\mu_z = 0$. The remaining parameters are the same as in Fig. 1. Temperature is measured in units of ε/k_B



Thus far we considered zero temperature, $T = 0$. The effect of increasing temperature is depicted in Fig. 4. Firstly, we observe that if temperature increases the phase does not vanish for $\theta \rightarrow \pi$ while it tends to zero for $\theta \rightarrow 0$. Secondly, the main properties remain similar: In all presented cases a minimum and a maximum exist. However, the maximum diminishes with increasing temperature.

5 Concluding remarks

No realistic physical quantum system is in perfect isolation from its environment. At best one can achieve a weak coupling between the system and the environment. In this weak coupling regime it is possible to extract the reduced dynamics of the open quantum system in a mathematically satisfactory and controlled way by using a Markovian reduced dynamics following the Davies scheme. In this work we have analyzed the geometric phase of a qubit in the presence of a weak coupling to a bosonic environment. We have investigated the relation between the geometric phase Φ and the mechanism for decoherence of the qubit for either the case of pure dephasing with $\mu_x = 0$ or in presence of dissipative energy relaxation, i.e. $\mu_x \neq 0$. The latter situation allows for a significant variation of the emerging geometric phase upon varying

the coupling strength μ_x . A variation of the pure dephasing coupling, i.e. $\mu_z \neq 0$ with $\mu_x = 0$, between qubit and environment barely affects the geometric phase. This feature is distinct from other set-ups, such as the emergence of quantum entanglement in open systems, where this dephasing-coupling mechanism can play a dominant or a similar role as an energy relaxation-coupling.

Nowadays, the geometric phase plays a crucial role in a variety of physical problems and has observable consequences in a wide range of systems. Under various aspects, this concept occurs in geometry, astronomy, classical mechanics, and quantum theory. The impressive recent progress in nanotechnology and experimental techniques allows one to test the fundamentals of quantum dynamics and details of interactions modeled by Hamiltonians. The geometric phase is not a quantum mechanical observable, i.e. it is not represented by a Hermitian operator. However, it can be experimentally measured, cf. Refs. [61–63]. It can be used to encode information on systems. E.g. it has been proposed as an order parameter for quantum phase transitions [64]. The results obtained here suggest that one can also exploit the geometric phase as a quantifier characterizing a nature of the system-environment coupling. Indeed, three features of the geometric phase $\Phi = \Phi(\mathcal{T})$ allow one to distinguish the character of qubit-environment coupling (i.e. pure dephasing vs. dissipation): (i) rotation symmetry around some points on the $\theta - \Phi$ plane or equivalently antisymmetric dependence of Φ upon θ about some points, (ii) non-monotonic behavior of Φ with respect to θ and (iii) the behavior of Φ for $\theta \rightarrow 0$ (i.e. for the qubit prepared in the excited state). We have verified that all these three features are manifest also at times $t = n\mathcal{T}$ ($n = 2, 3, 4$) for the measurable quantifier $\Phi = \Phi(n\mathcal{T})$. This feature of the geometric phase Φ thus presents an additional suitable tool in exploring characteristics of open system interactions at the quantum scale.

Acknowledgment Work supported by the Polish Ministry of Science and Higher Education under the grant N 202 131 32/3786, the German Excellence Initiative via the “Nanosystems Initiative Munich (NIM)” and by the DFG-SFB 631.

References

1. Nielsen, M.A., Chuang, I.L.: Quantum Computation and Quantum Information. Cambridge University Press, Cambridge (2000)
2. Lidar, D.A., Whalley, K.B.: Irreversible quantum dynamics. Lecture Notes in Physics, vol. 622, 83. Springer, Berlin (2006)
3. Alicki, R.: *ibid*, **121**
4. Kohler, S., Hänggi, P.: Improving the purity of one- and two-qubit gates. Fortschr. Physik **54**, 804–819 (2006)
5. Zanardi, P., Rasetti, M.: Holonomic quantum computation. Phys. Lett. A **264**, 94–99 (1999)
6. Nayak, C. et al.: Non-Abelian anyons and topological quantum computation. Rev. Mod. Phys. **80**, 1083–1159 (2008)
7. Jones, J.A., Vedral, V., Ekert, A., Castagnoli, G.: Geometric quantum computation using nuclear magnetic resonance. Nature (London) **403**, 869–871 (2000)
8. Sarandy, M.S., Lidar, D.A.: Adiabatic quantum computation in open systems. Phys. Rev. Lett. **95**, 250503–250507 (2005)
9. Berry, M.V.: Quantal phase factors accompanying adiabatic changes. Proc. R. Soc. London Ser. A **329**, 45–57 (1984)

10. Wilczek, F., Zee, A.: Appearance of gauge structure in simple dynamical systems. *Phys. Rev. Lett.* **52**, 2111–2114 (1984)
11. Duan, L.-M., Cirac, J.I., Zoller, P.: Geometric manipulation of trapped ions for quantum computation. *Science* **292**, 1695–1697 (2001)
12. Recati, A., Calarco, T., Zanardi, P., Cirac, J.I., Zoller, P.: Holonomic quantum computation with neutral atoms. *Phys. Rev. A* **66**, 032309–032322 (2002)
13. Yin, S., Tong, M.D.: Geometric phase of a quantum dot system in nonunitary evolution. *Phys. Rev. A* **79**, 044303–044307 (2009)
14. Falci, G., Fazio, R., Palma, G.M., Siewert, J., Vedral, V.: Detection of geometric phases in superconducting nanocircuits. *Nature* **407**, 355–358 (2000)
15. Faoro, L., Siewert, J., Fazio, R.: Non-abelian phases, charge pumping and holonomic computation with Josephson junctions. *J. Phys. Soc. Jpn.* **72**, 3–4 (2003)
16. Parodi, D., Sasseti, M., Solinas, P., Zanardi, P., Zangh, N.: Fidelity optimization for holonomic quantum gates in dissipative environments. *Phys. Rev. A* **73**, 052304–052309 (2006)
17. Parodi, D., Sasseti, M., Solinas, P., Zangh, N.: Environmental noise reduction for holonomic quantum gates. *Phys. Rev. A* **76**, 012337–012343 (2007)
18. Uhlmann, A.: The transition probability in the state space of a $*$ -algebra. *Rep. Math. Phys.* **9**, 273–279 (1976)
19. Bassi, A., Ippoliti, E.: Geometric phase for open quantum systems and stochastic unravellings. *Phys. Rev. A* **73**, 062104–062111 (2006)
20. Burić, N., Radonjić, M.: Uniquely defined geometric phase of an open system. *Phys. Rev. A* **80**, 014101–014105 (2009)
21. Sjöqvist, E. et al.: Geometric phases for mixed states in interferometry. *Phys. Rev. Lett.* **85**, 2845–2849 (2000)
22. Bhandari, R.: Singularities of the mixed state phase. *Phys. Rev. Lett.* **89**, 268901 (2002)
23. Sjöqvist, E.: Quantal interferometry with dissipative internal motion. *Phys. Rev. A* **70**, 052109–052115 (2004)
24. Bhandari, R.: Polarization of light and topological phases. *Phys. Rep.* **281**, 1–64 (1997)
25. Du, J. et al.: An experimental observation of geometric phases for mixed states using NMR interferometry. *Phys. Rev. Lett.* **91**, 100403–100407 (2003)
26. Mukunda, N., Simon, R.: Quantum kinematic approach to the geometric phase I. General formalism. *Ann. Phys.* **228**, 205–268 (1993)
27. Tong, D.M., Sjöqvist, E., Kwek, L.C., Oh, C.H.: Kinematic approach to the mixed state geometric phase in nonunitary evolution. *Phys. Rev. Lett.* **93**, 080405–080409 (2004)
28. Carollo, A., Fuentes-Guridi, I., Frana Santos, M., Vedral, V.: Geometric phase in open systems. *Phys. Rev. Lett.* **90**, 160402–160406 (2003)
29. Ericsson, M., Sjöqvist, E., Brännlund, J., Oi, D.K., Pati, A.K.: Generalization of the geometric phase to completely positive maps. *Phys. Rev. A* **67**, 020101–020105 (2003)
30. Marzlin, K.-P., Ghose, S., Sanders, B.C.: Geometric phase distributions for open quantum systems. *Phys. Rev. Lett.* **93**, 260402–260406 (2004)
31. Whitney, R.S., Makhlin, Y., Shnirman, A., Gefen, Y.: Geometric nature of the environment-induced Berry phase and geometric dephasing. *Phys. Rev. Lett.* **94**, 070407–070411 (2005)
32. Sarandy, M.S., Duzzioni, E.I., Moussa, M.H.Y.: Dynamical invariants and nonadiabatic geometric phases in open quantum systems. *Phys. Rev. A* **76**, 052112–052121 (2007)
33. Huang, X.L., Yi, X.X.: Non-Markovian effects on the geometric phase. *Europhys. Lett.* **82**, 50001–50007 (2008)
34. Banerjee, S., Srikanth, R.: Geometric phase of a qubit interacting with a squeezed-thermal bath. *Eur. Phys. J. D* **46**, 335–344 (2008)
35. Fujikawa, K., Hu, M.-G.: Geometric phase of a two-level system in a dissipative environment. *Phys. Rev. A* **79**, 052107–052114 (2009)
36. Wang, Z.S., Liu, G.Q., Ji, Y.H.: Noncyclic geometric quantum computation in a nuclear-magnetic-resonance system. *Phys. Rev. A* **79**, 054301–054305 (2009)
37. Singh, K. et al.: Geometric phases for nondegenerate and degenerate mixed states. *Phys. Rev. A* **67**, 032106–032115 (2003)
38. Hänggi, P., Ingold, G.L.: Fundamental aspects of quantum Brownian motion. *Chaos* **15**, 026105–026115 (2005)

39. Alicki, R., Fannes, M., Pogorzelska, M.: Quantum generalized subsystems. *Phys. Rev. A* **79**, 052111–052120 (2009)
40. Łuczka, J.: Spin in contact with thermostat: Exact reduced dynamics. *Physica A* **167**, 919–934 (1990)
41. Alicki, R.: Pure decoherence in quantum systems. *Open Sys. & Inf. Dyn.* **11**, 53–61 (2004)
42. Romero, K.M.F., Talkner, P., Hänggi, P.: Is the dynamics of open quantum systems always linear? *Phys. Rev. A* **69**, 052109–052117 (2004)
43. Dajka, J., Mierzejewski, M., Łuczka, J.: Fidelity of asymmetric dephasing channels. *Phys. Rev. A* **79**, 012104–012111 (2009)
44. Doll, R., Wubs, M., Hänggi, P., Kohler, S.: Limitation of entanglement due to spatial qubit separation. *Europhys. Lett.* **76**, 547–553 (2006)
45. Doll, R., Wubs, M., Hänggi, P., Kohler, S.: Incomplete pure dephasing of N-qubit entangled W states. *Phys. Rev. B* **76**, 045317–045331 (2007)
46. Dajka, J., Mierzejewski, M., Łuczka, J.: Entanglement persistence in contact with the environment: exact results. *J. Phys. A: Math. Theor.* **40**, F879–F886 (2007)
47. Dajka, J., Łuczka, J.: Origination and survival of qudit-qudit entanglement in open systems. *Phys. Rev. A* **77**, 062303–062310 (2008)
48. Doll, R., Hänggi, P., Kohler, S., Wubs, M.: Fast initial qubit decoherence and the influence of substrate dimensions on error correction rates. *Eur. Phys. J. B* **68**, 523–527 (2009)
49. Yi, X.X., Wang, L.C., Wang, W.: Geometric phase in dephasing systems. *Phys. Rev. A* **71**, 044101–044105 (2005)
50. Yi, X.X., Tong, D.M., Wang, L.C., Kwek, L.C., Oh, C.H.: Geometric phase in open systems: beyond the Markov approximation and weak-coupling limit. *Phys. Rev. A* **73**, 052103–052109 (2006)
51. Dajka, J., Mierzejewski, M., Łuczka, J.: Geometric phase of a qubit in dephasing environment. *J. Phys. A Math. Theor.* **41**, F012001–F012008 (2008)
52. Dajka, J., Łuczka, J.: Bifurcations of the geometric phase of a qubit asymmetrically coupled to the environment. *J. Phys. A: Math. Theor.* **41**, F442001–F442009 (2008)
53. Davies, E.B.: Markovian master equations. *Comm. Math. Phys.* **39**, 91–110 (1974)
54. Dümcke, R., Spohn, H.: The proper form of the generator in the weak coupling limit. *Z. Physik B* **34**, 419–422 (1979)
55. Łuczka, J.: On Markovian kinetic equations: Zubarev's nonequilibrium statistical operator approach. *Physica A* **149**, 245–266 (1988)
56. Lendi, K., van Wonderen, A.J.: Davies theory for reservoir-induced entanglement in a bipartite system. *J. Phys. A Math. Theor.* **40**, 279–288 (2007)
57. Schuster, D.I. et al.: Resolving photon number states in a superconducting circuit. *Nature* **445**, 515–518 (2007)
58. Alicki, R., Lendi, K.: *Quantum dynamical semigroups and applications*. Springer, Berlin (1987)
59. Aharonov, Y., Anandan, J.: Phase change during a cyclic quantum evolution. *Phys. Rev. Lett.* **58**, 1593–1597 (1987)
60. Chruściński, D., Jamiołkowski, A.: *Geometric phases in classical and quantum mechanics*. Birkhäuser, Boston (2004)
61. Leek, P.J. et al.: Observation of Berry's phase in a solid-state qubit. *Science* **318**, 1889–1892 (2007)
62. Möttönen, M. et al.: Experimental determination of the Berry phase in a superconducting charge pump. *Phys. Rev. Lett.* **100**, 177201–177205 (2008)
63. Fillipp, S. et al.: Experimental demonstration of the stability of Berry's phase for a spin-1/2 particle. *Phys. Rev. Lett.* **102**, 030404–030408 (2009)
64. Nesterov, A.I., Ovchinnikov, S.G.: Geometric phases and quantum phase transitions in open systems. *Phys. Rev. E* **78**, 015202–015206 (2008)